

syngenta

FIFRA

## Section 2(ee) Recommendation

ACCEPTED  
FOR REGISTRATION

1/28/04

New York State Department  
of Environmental Conservation  
Division of Solid &  
Hazardous Materials

Date: January 27, 2004

Products: Medallion® Fungicide, EPA Reg. No. 100-769  
Daconil Ultrex®, EPA Reg. No. 50534-202-100

Use: Application rates of 0.25 to 0.5 oz/1000 sq ft of Medallion applied alone in alternation with other fungicides or as a tank mixture with 3.25 oz/1000 sq ft of Daconil Ultrex for the control of anthracnose in annual bluegrass and creeping bentgrass.

States: New York (except Nassau & Suffolk counties)

This recommendation is made under the authority of FIFRA Section 2(ee) and recommends a preventive application of Medallion at 0.25 to 0.5 oz./1000 sq ft either alone or in combination with Daconil Ultrex at 3.25 oz./1000 sq ft on for the control of anthracnose in New York.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### Control of Anthracnose in Poa annua and Creeping Bentgrass

#### General Information

Medallion and Daconil Ultrex are protectant fungicides that provide broad spectrum disease control in turfgrass. Daconil Ultrex is one of only a few fungicides labeled for the control of anthracnose. In recent testing, Medallion applied alone provided good preventive control of anthracnose. Medallion, when applied in a tank mixture with Daconil Ultrex, may improve the control of anthracnose and other diseases. Alternating applications or tank-mixing different fungicide chemistries provides optimal disease control and minimizes the potential risk of resistance development.

### **Directions For Use**

This FIFRA 2(ee) recommendation allows the use of Medallion alone or Medallion plus Daconil Ultrex as a tank-mixture in a preventive disease control program targeted at anthracnose. Follow the label directions for use of Medallion and Daconil Ultrex to control labeled turfgrass diseases. The first application of a fungicide should be initiated 3-4 weeks prior to disease development. Use historical data to determine the typical anthracnose disease development start date. Application intervals should not be any greater than 14 days. Limit the number of Medallion applications to 6 applications at 0.25 oz/1000 sq ft, 4 applications at 0.33 oz/1000 sq ft, or 3 applications at 0.5 oz/1000 sq ft not to exceed a total of 1.5 oz of Medallion per 1000 ft<sup>2</sup> per year.

### **Restrictions:**

- (1) The user must refer to the federally approved labels for the above noted products and read and follow all directions for use, restrictions, and precautions.
- (2) Do not apply more than 1.5 oz/1000 ft<sup>2</sup>/year of Medallion on turfgrass.
- (3) The user should have this recommendation in its possession at the time of use.
- (4) This is **NOT** a permitted use in Nassau and Suffolk Counties, New York.

This recommendation for use of Medallion in tank mixture with Daconil Ultrex for control of anthracnose is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Syngenta Crop Protection, Inc.  
Greensboro, NC 27419

Medallion<sup>®</sup>, Daconil Ultrex<sup>®</sup>, and the Syngenta logo are trademarks of a Syngenta Group Company.

©2003 Syngenta.

Label code: NY0769010AC0104