

# New York State Department of Environmental Conservation

## Division of Solid & Hazardous Materials


Bureau of Pesticides Management

625 Broadway, Albany, New York 12233-7254

Phone: 518-402-8788 FAX: 518-402-9024

Website: [www.dec.ny.gov](http://www.dec.ny.gov)

E-Mail: [ppr@gw.dec.state.ny.us](mailto:ppr@gw.dec.state.ny.us)



Alexander B. Grannis  
Commissioner

June 15, 2007

**CERTIFIED MAIL**  
**RETURN RECEIPT REQUESTED**

Ms. Miriam Carr  
E. I. Dupont De Nemours & Company  
Stine-Haskell Research Center, Suite #300/427  
P.O. Box 30  
Newark, Delaware 19714-0030

Dear Ms. Carr:

**Re: Registration of Provaunt Insecticide (EPA Reg. No. 352-716) Containing the Active Ingredient Indoxacarb. Chemical Code: 067710.**

The New York State Department of Environmental Conservation (Department) has completed its technical review of the above product application received on October 10, 2006. The application was submitted as a routine product application which would contain active ingredients and use patterns already registered for use in New York State. Indoxacarb is currently registered in New York State for agricultural use and for indoor use to control ants and cockroaches. Provaunt Insecticide is labeled for use on turf including golf courses and landscape ornamentals. The application was determined to be a major change in labeled use pattern (MCL) due to the addition of the turf and ornamental use pattern. Dupont provided the additional data on December 20, 2006 regarding human exposure to treated turf and the Department determined that this additional information along with the existing data on file for the product DuPont Avaunt Insecticide (EPA Reg. No. 352-597) was sufficient to determine the application complete for technical review on February 16, 2007.

Provaunt Insecticide is formulated as a water dispersible granule containing 30% indoxacarb and is labeled for use on turf including landscape and recreational turf (including golf courses) and landscape ornamentals. The maximum application rate is 0.45 lb. of indoxacarb per acre per year as a broadcast application on turf grass. The agricultural use of the same formulation known as DuPont Avaunt Insecticide (EPA Reg. No. 352-597) is applied as a foliar application on certain crops at a maximum annual use rate of 0.44 lb. ai/acre/yr. Indoxacarb belongs to the oxadiazine chemical class.

Pursuant to the review time frame specified in ECL §33-0704.2, a registration decision date of July 16, 2007 was established. The Department has conducted the following technical review for impacts to human health as a result of potential increased exposure to the active ingredient indoxacarb when used on turf and ornamentals. The following review summary is provided below.

**Human Health Risk Assessment:**

According to the registrant (per correspondence from S. Theodorakis to S. Jackling, December 20, 2006), "...the formulation for DuPont Provaunt Insecticide is the same formulation as the currently registered crop use product DuPont Avaunt Insecticide (EPA Reg. No. 352-597)." The New York State Department of Health (DOH) previously reviewed Avaunt Insecticide which was subsequently registered in New York State on March 21, 2002. Both Avaunt and indoxacarb overall were not very acutely toxic in laboratory animal studies. Indoxacarb also did not cause reproductive or carcinogenic effects and was negative in a number of genotoxicity studies. Indoxacarb caused some developmental effects in laboratory animals, but only at doses that also caused maternal effects and in studies that identified no-observed-effect levels (NOELs). In chronic feeding animal studies, indoxacarb caused some toxicity, including decreased body weight, body weight gain and food consumption in male and female rats as well as decreased hematocrit, hemoglobin and red blood cells in females. Using data from the chronic rat feeding study, the United States Environmental Protection Agency (USEPA) Office of Pesticide Programs established a reference dose for indoxacarb of 0.02 milligrams per kilogram body weight per day (mg/kg/day) based on a NOEL of 2.1 mg/kg/day (for hematological effects) and an uncertainty factor of 100. A current search of the toxicological literature did not find any significant new information on the toxicity of indoxacarb.

The registrant submitted a USEPA risk assessment for occupational, residential and recreational exposures to indoxacarb resulting from the labeled uses of the DuPont Provaunt product. For workers, the estimated range of dermal and inhalation margins of exposure (MOEs) were 800 to 31,000, and 800 to 55,000, respectively. The dermal MOEs were determined by comparing the estimated dermal exposures to a NOEL of 50 mg/kg/day obtained from a 28-day dermal toxicity study conducted on rats (decreases in body weights and body weight gains, occurrence of hematological effects), whereas the inhalation MOEs were determined by comparing the estimated inhalation exposures to a NOEL of 6 mg/kg/day obtained from a 28-day rat inhalation toxicity study (increase in spleen weights, hematological changes). In estimating these MOEs, it was assumed that workers wore personal protective equipment (long-sleeved shirt, long pants, shoes plus socks, and chemical-resistant gloves) as is required by the product label. Generally, the USEPA considers MOEs of 100-fold or greater to provide adequate worker protection. For post-application exposures to indoxacarb, the estimated dermal MOEs for golf course workers, adult and child golfers were 110, 4,166 and 2,500, respectively. For residential post-application dermal exposures, the estimated MOEs for children and adults were 166 and 277, respectively. Finally, the estimated MOEs for post-application oral exposures of children ranged from 166 (hand to mouth from contacting treated turf) to about 133,000 (incidental ingestion of treated soil). For determining the oral MOEs, estimated exposures were compared to a NOEL of 2 mg/kg/day obtained from a subchronic rat feeding study (decreases in body weights and body weight gains).

There are no chemical specific federal or New York State drinking water/groundwater standards for indoxacarb. Based on its chemical structure, indoxacarb falls under the 50 microgram per liter New York State drinking water standard for an "unspecified organic contaminant" (10 NYCRR Part 5, Public Water Systems).

The available information on indoxacarb and the formulated product DuPont Provaunt Insecticide indicates that overall they were not very acutely toxic in laboratory animal studies. Indoxacarb also did not cause reproductive or carcinogenic effects, and only caused

developmental effects at maternally toxic doses. Although data from subchronic and chronic animal feeding studies on indoxacarb showed that this chemical has the potential to cause some toxicity, the expected exposures from using the DuPont Provaunt Insecticide product should not pose a significant risk to workers or the general public. Given the above, we do not object to the registration of DuPont Provaunt Insecticide in New York State.

**Product Registration Summary:**

The Department will register Dupont Provaunt Insecticide (EPA Reg. No. 352-716) for general use in New York State. The initial registration of Avaunt Insecticide required a New York State specific labeling statement to prohibit sale and use of the product on Long Island, New York due to the lack of data available to thoroughly review impacts to groundwater. Dupont submitted data and an application to this Department on October 26, 2005 for the purpose of removing the statement "Not for Use or Sale in Suffolk and Nassau Counties, Long Island, New York" from the DuPont Avaunt Insecticide label. This Department conducted an environmental fate assessment and concluded that given the short half-lives, the low application rates, and the Kocs, LEACHP modeling projected no impact to groundwater from the aerobic metabolism transformation products of the precursor indoxacarb. Therefore, DuPont's request to remove the Long Island prohibition was approved for Avaunt Insecticide and subsequently will not be required on the Provaunt Insecticide label.

Enclosed for your record, is a copy of the stamped accepted labeling and the Certificate of Registration for DuPont Provaunt Insecticide (EPA Reg. No. 352-716). Please note that a proposal by E. I. DuPont De Nemours & Company or any other registrant, to register a product that contains indoxacarb, and whose labeled uses are likely to increase the potential for significant impact to humans, nontarget organisms, or the environment, would constitute a MCL. Such an application must be accompanied by a new application fee and meet the requirements listed in Appendix 1.B. of "New York State Pesticide Product Registration Procedures" (September 2005). Such information as well as forms can be accessed at our website as listed in our letterhead.

Please be aware that any unregistered product may **not** be sold, offered for sale, distributed, or used in New York State.

Please contact our Pesticide Product Registration Section, at (518) 402-8768, if you have any questions.

Sincerely,

*Maureen P Serafini*

Maureen P. Serafini  
Director  
Bureau of Pesticides Management

Enclosure

cc: w/enc. - N. Kim/D. Luttinger - NYS Dept. of Health  
R. Mungari - NYS Dept. of Ag. & Markets  
W. Smith - Cornell University, PSUR

# DuPont™ Provaunt™

## INSECTICIDE

### Professional Products

GROUP	22	INSECTICIDE
-------	----	-------------

A water dispersible granular insecticide for the control of lepidopterous larvae (including armyworms, cutworms, sod webworms, bagworms, fall webworms, gypsy moth caterpillars, tent caterpillars, tussock moth caterpillars and yellownecked caterpillars) and certain other pests infesting landscape and recreational (including golf courses) turfgrass and landscape ornamentals.

<i>Active Ingredient</i>	<i>By Weight</i>
--------------------------	------------------


Indoxacarb*	
(S)-methyl 7-chloro-2,5-dihydro-2-[[methoxy-carbonyl][4(trifluoromethoxy)phenyl]amino]-carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a-(3H)-carboxylate	30.0%
<i>Other ingredients</i>	70.0%
<b>TOTAL</b>	<b>100.0%</b>

\*Indoxacarb belongs to the oxadiazine chemical class.

EPA Reg. No. 352-716

NET CONTENTS: \_\_\_\_\_

Establishment No.:



### PRECAUTIONARY STATEMENTS

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

### FIRST AID

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or a doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Remove contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information. For information on the use of this product call DuPont Professional Products at 1-888-6-DUPONT (1-888-638-7668).



*The miracles of science™*

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution!: Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing (dust, vapor or spray mist). Remove contaminated clothing and wash clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all > 14 mls.
- Shoes plus socks

After the product has been diluted in accordance with label directions for use, shirt, pants, socks, shoes and waterproof gloves are sufficient Personal Protective Equipment. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

**USER SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production.
- Do not formulate this product into other end-use products without written permission of DuPont Professional Products
- Do not apply this product through any type of irrigation system.
- Do not apply this product in commercial nurseries and greenhouses or on sod farms.
- Do not apply more than 24 ounces (equivalent to 0.45 lb of active ingredient) of product per acre per season in broadcast applications to turfgrass.
- Wait a minimum of 7 days to retreat.
- Keep people and pets away from treated areas until treatment has dried.

DuPont™ PROVAUNT™ insecticide should be used only in accordance with recommendations on this label or in separate DuPont supplemental labeling that may be made temporarily available through local distributors, as a result of new EPA approvals. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont. User assumes all risks associated with such non-recommended use.

### GENERAL INFORMATION

PROVAUNT™ insecticide is a water dispersible granular that may be applied as a foliar or broadcast spray to control lepidopterous larvae (such as armyworms, cutworms, sod webworms, bagworms, fall webworms, gypsy moth caterpillars, tent caterpillars, tussock moth caterpillars, yellownecked caterpillars) and certain other pests on landscape and recreational (including golf course) turfgrass and landscape ornamentals including trees, shrubs, foliage plants, flowers and non-bearing fruit and nut trees, which will not produce a raw agricultural commodity during the season of application. PROVAUNT™ insecticide may only be used on plants or turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries, ornamental gardens, parks, playgrounds, schools, golf courses (tee box areas, roughs, fairways, greens, collars etc.), athletic fields and other landscaped areas.

PROVAUNT™ insecticide is in GROUP 22 of the EPA's Insecticide and Acaricide Groups Based on Target Site of Action (EPA PR Notice 2001-5) and may be used in rotational resistance management programs. PROVAUNT™ insecticide must be diluted with water before application.

## INTEGRATED PEST MANAGEMENT (IPM) PROGRAMS

DuPont™ PROVAUNT™ insecticide is recommended for IPM programs on turf and landscape ornamentals because it does not directly impact natural arthropod predator and parasitoid populations including ladybird beetles, lacewings, minute pirate bugs and predatory mites. The feeding behavior of predatory beneficial arthropods will aid in extending natural control of other insect and mite pests and will reduce the possibility of secondary pest outbreaks. PROVAUNT™ insecticide will reduce the target pest species that may serve as a food source for beneficial arthropods. If PROVAUNT™ insecticide is tank-mixed with an insect control product that negatively impacts beneficial arthropods, then the full benefit of PROVAUNT™ insecticide to the IPM program may not be realized.

### APPLICATION TIMING

PROVAUNT™ insecticide applications should target eggs and small instar lepidopterous larvae. Use sufficient water to obtain thorough, uniform coverage. Repeat applications may be made after 7 days, if necessary.

### APPLICATION EQUIPMENT PREPARATION

Application equipment must be clean and free of previous pesticide deposits before mixing PROVAUNT™ insecticide.

#### Mixing Directions:

1. Use clean, well maintained application equipment.
2. Fill sprayer tank 1/4 to 1/2 full of water.
3. Then add DuPont™ PROVAUNT™ insecticide directly to the sprayer tank.
5. Mix thoroughly to fully disperse the insecticide and continue agitation to keep the insecticide in suspension. Use mechanical or hydraulic agitation. Do not use air agitation.
6. The mixture should not be stored in the spray tank overnight.

#### Tank-mixtures:

PROVAUNT™ insecticide may be tank-mixed with other pesticides. When tank-mixing PROVAUNT™ insecticide with other pesticides, observe all precautions and limitations on each separate product label. No label application rates may be exceeded. PROVAUNT™ insecticide cannot be mixed with any product containing a label prohibition against such mixing. The physical compatibility of PROVAUNT™ insecticide may vary with different sources of pesticide products and local cultural practices. Any tank-mixture that has not been previously tested should be prepared on a small scale (pint or quart jar) using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture.

#### Tank-mixing Sequence:

Add different formulation types in the sequence indicated below. Allow time for complete mixing and dispersion after the addition of each product.

1. Water soluble bags
2. PROVAUNT™ and other water dispersible granules
3. Wettable powders

4. Water-based suspension concentrates
5. Water soluble concentrates
6. PROVAUNT™ insecticide and other oil-based suspension concentrates
7. Emulsifiable concentrates
8. Adjuvants, surfactants, oils
9. Soluble fertilizers
10. Drift retardants

### APPLICATION EQUIPMENT CLEANING

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all application equipment to reduce the risk of forming hardened deposits that might become difficult to remove. Drain application equipment. Thoroughly rinse application equipment and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean equipment near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

### RESISTANCE MANAGEMENT

Some insects are known to develop resistance to products used repeatedly for control. When this occurs, the recommended dosages fail to suppress the pest population below threshold levels. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of active classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local DuPont Professional Products representative, Cooperative Extension Service specialist or pest control advisor for the latest information on resistance management in your area.

### APPLICATION RATES FOR LAWNS, GOLF COURSES AND OTHER RECREATIONAL-TURFGRASS AREAS

To ensure optimum control of lepidopterous larvae, delay watering (irrigation) and mowing for 24 hours after application. If the grass area is being maintained at a mowing height of greater than 1 inch, then the higher application rate (4 ounces per acre) may be required during periods of high pest pressure. For maximum residual effectiveness, PROVAUNT™ insecticide may be applied at up to 12 ounces per acre (0.275 ounces per 1,000 square feet = 0.225 lb A/A).

Target Pest	Product per Acre	Product per 1,000 Square Feet	lb A/A
Caterpillars (including armyworms, cutworms and sod webworms)	2 to 4 Ounces	0.046 to 0.092 Ounces	0.0375 to 0.075
European Crane Fly Larvae	6 to 12 Ounces	0.138 to 0.275 Ounces	0.1125 to 0.225
Grasshoppers	12 Ounces	0.275 Ounces	0.225

## APPLICATION RATES FOR ORNAMENTAL PLANTS (EXTERIOR LANDSCAPES)

Apply the desired application rate of DuPont™ PROVAUNT™ insecticide as a full coverage foliar spray. Repeat treatment as necessary to achieve control using higher application rates as pest pressure and foliage area increases. Repeat applications must be limited to no more than once per seven days. Certain plant species or cultivars may be sensitive to the final spray solution. If local experience is not available, then a small number of plants should be treated and observed for phytotoxicity for at least one week before making application to the entire planting.

Target Pests	Product per 100 Gallons	lb AI per 100 Gallons	PPM	Percent AI
Leaf-feeding caterpillars (such as bagworms, gypsy moth caterpillars, tent caterpillars [including eastern tent caterpillar], tussock moth caterpillars [including whitemarked tussock moth], webworms [including fall webworm] and yellownecked caterpillars)	1.25 to 2.5 Ounces	0.023 to 0.046	28.1 to 56.2	0.0028 to 0.0056
European pine sawfly	2.5 to 5.0 Ounces	0.046 to 0.092	56.2 to 112.4	0.0056 to 0.01124
Potato leafhopper	5.0 Ounces	0.092	112.4	0.01124

## APPLICATION DILUTION CHART

Application Volume (Gallons per 1,000 Square Feet)	Application Rate			Ounces of DuPont™ PROVAUNT™ diluted to these volumes of finished spray			
	Ounces per Acre	Ounces per 1,000 Square Feet	lb A.I. per Acre	1 gallon	5 gallons	10 gallons	100 gallons
1	2	0.046	0.0375	0.046	0.23	0.46	4.6
	3	0.069	0.05625	0.069	0.345	0.69	6.9
	4	0.092	0.075	0.092	0.46	0.92	9.2
	6	0.138	0.1125	0.138	0.69	1.38	13.8
	12	0.275	0.225	0.275	1.38	2.76	27.6
2	2	0.046	0.0375	0.023	0.115	0.23	2.3
	3	0.069	0.05625	0.035	0.173	0.345	3.45
	4	0.092	0.075	0.046	0.23	0.46	4.6
	6	0.138	0.1125	0.069	0.345	0.69	6.9
	12	0.275	0.225	0.138	0.69	1.38	13.8
3	2	0.046	0.0375	0.015	0.077	0.154	1.54
	3	0.069	0.05625	0.023	0.116	0.231	2.31
	4	0.092	0.075	0.031	0.154	0.308	3.08
	6	0.138	0.1125	0.046	0.231	0.462	4.62
	12	0.275	0.225	0.092	0.462	0.924	9.24
4	2	0.046	0.0375	0.012	0.058	0.116	1.16
	3	0.069	0.05625	0.017	0.087	0.174	1.74
	4	0.092	0.075	0.023	0.116	0.232	2.32
	6	0.138	0.1125	0.035	0.174	0.348	3.48
	12	0.275	0.225	0.070	0.348	0.696	6.96

- To convert from ounces to grams, multiply by 28.35.
- 1 ounce = 28.35 grams.
- Do not use household utensils to measure PROVAUNT™ insecticide

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**NOTICE TO BUYER**— Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™ and PROVAUNT™ are trademarks of E.I. duPont de Nemours and Company.

SL - 1113A 081406 07-10-06

## LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont or Seller. These risks can cause: ineffectiveness of the product or other unintended consequences. DuPont does not agree to be an insurer of these risks. **TO THE FULLEST EXTENT PERMITTED BY LAW, WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHER LEGAL THEORY) RESULTING FROM THE STORAGE, USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

DuPont or its distributor must have prompt notice of any claim so that an immediate investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify DuPont or a DuPont distributor of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy. This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT (1-888-638-7668)

Internet address: <http://www.proproducts.dupont.com/>

© 2006 E.I. du Pont de Nemours and Company, Wilmington, Delaware 19898

All rights reserved.