



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014, 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

STRATEGO® Fungicide

**For Use on Soybean to control Rust
 (*Phakopsora pachyrhizi*)**

**SECTION 18 QUARANTINE EXEMPTION
 File Symbol: 04-NY-20**

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEW YORK
 UNDER SECTION 18 QUARANTINE EXEMPTION**

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for STRATEGO® Fungicide (EPA Registration Number 264-779).

Any adverse effects resulting from the use of STRATEGO® Fungicide under this quarantine exemption must be immediately reported to the New York State Department of Environmental Conservation.

EFFECTIVE PERIOD:

This labeling becomes effective 01/07/05 and expires on 12/01/07.

A maximum of 2 total applications using approved Section 18 products collectively will be made under this soybean rust Section 18.

DIRECTIONS FOR USE

A maximum of 140,000 acres of soybeans may be treated under this exemption.

RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF STRATEGO® FUNGICIDE
Soybean	Rust (<i>Phakopsora pachyrhizi</i>)	5.5 to 10 fl oz per acre

Use Directions: Begin broadcast, foliar applications of STRATEGO® preventatively when conditions are favorable for disease development. Continue application programs as needed for disease control using a 10 to 21-day spray interval. Applications of STRATEGO® may be made up to the R6 (full seed) soybean growth stage. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. Do not apply STRATEGO® in combination with organosilicate surfactants or crop injury may occur. Use of adjuvants may enhance performance of STRATEGO®. If utilized, apply the lowest labeled rate of the adjuvant to enhance disease control. STRATEGO® should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment. Chemical-resistant gloves must be worn for mixers/loaders. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 2 applications per season. Do not apply more than 20 fl. oz/A per use season. Do not enter treated fields for 24 hours after application.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)
 or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

