

This labeling is effective for distribution and use of this product to control soybean rust on soybeans grown in a limited number of states until April 19, 2009, as authorized pursuant to the provisions of section 18 of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended.

# Caramba™


fungicide

**Emergency Exemption Use Directions for use in Soybean for the control of Asian Soybean Rust (*Phakopsora pachyrhizi*)**

EPA File Symbol No.: 07-NY-05

Caramba™ fungicide contains 0.75 pounds of metconazole per gallon

**OBSERVE ALL FIRST AID, PRECAUTIONARY STATEMENTS, AND DIRECTIONS FOR USE ON THE CARAMBA™ FUNGICIDE CONTAINER LABEL BEFORE USING.**



BASF Corporation  
Agricultural Products  
26 Davis Drive  
Research Triangle Park, NC 27709

 **BASF**  
The Chemical Company

## DIRECTIONS FOR USE

### Caramba™ fungicide

#### For Use in Soybean for the control of Asian Soybean Rust caused by *Phakopsora pachyrhizi*

The use of Caramba™ fungicide for control of soybean rust in soybeans in a limited number of states as authorized under the provisions of section 18 of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is effective until April 19, 2009. In accordance with 40 CFR §152.30(e), an unregistered pesticide may be distributed or sold in accordance with the terms of an emergency exemption under FIFRA section 18, if the product is properly labeled in accordance with 40 CFR §156. Any unused, unregistered product must be either returned to the manufacturer or distributor (unopened containers) or disposed of in accordance with Resource Conservation and Recovery Act (RCRA) regulations following the expiration of this authorization.

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

#### Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** discharge effluent containing this product into lakes streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

#### Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or

adversely modify their habitat is a violation of federal law. This pesticide is toxic to fish and aquatic invertebrates and must be used strictly in accordance with drift precautions on this label in order to minimize off-site exposures.

**DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitats. Notify State and/or Federal authorities and BASF immediately if you observe any adverse environmental effects due to use of this product.

To determine whether your county has endangered aquatic species, consult the County Bulletins at <http://www.epa.gov/espp/usa-map.htm>. Endangered Species Bulletins may also be obtained from extension offices or state pesticide agencies. If a bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered aquatic species occur in the area to be treated.

#### Restrictions and Limitations

- Applications may not be made within 30 days of harvest.
- **DO NOT** make more than 2 applications of metconazole-containing products per season.
- **DO NOT** feed soybean forage or hay to livestock (e.g., aspirated grain fractions, soybean hulls, soybean meal, or soybean seed).
- **Crop Rotation Restriction:** Any crop not specified on the label may be planted into treated areas as follows: for root crops and leafy vegetables, a plant-back interval (PBI) of 120 days must be observed. Wheat, corn, cotton, and rice may be planted 180 days after the last treatment. For all other crops, a 12-month PBI must be observed.
- **Chemigation: DO NOT** apply this product through any type of irrigation system.

- No aerial application in New York State except as permitted under FIFRA Section 24(c), Special Local Needs Registration.

## APPLICATION INSTRUCTIONS

**Caramba™** fungicide may be applied by ground sprayer or aerial equipment.

**Caramba** can be applied at a maximum rate of 9.6 fl oz per acre per application.

### Caramba Disease Specific Use Directions

Target Disease	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)	Restricted Entry Interval (REI)
Asian soybean rust* ( <i>Phakopsora pachyrhizi</i> )	8.2 to 9.6 fl oz per acre	2	19.2 fl oz per acre	30 days	12 hours

**Caramba** provides both preventative and curative activity against Asian soybean rust. For optimal disease control, make the initial application of **Caramba** between early flowering and pod set (R1 to R3 growth stage), or prior to disease development. If environmental conditions favor continued disease development, or if monitoring shows active disease symptoms, repeat application of **Caramba** 10-21 days after first application.

Use the higher rate and shorter spray interval when disease pressure is high.

Optimum disease control is achieved by utilizing the combination of **Caramba** and a preventative fungicide like **Headline®** fungicide. **Caramba** may be used at 6.1 fl oz per acre when tank mixed with a low rate of **Headline**.

Tank mixtures of **Caramba** with adjuvants or agricultural products containing adjuvants may cause crop response under hot dry conditions. If an adjuvant is needed, use only nonionic surfactants at their lowest labeled rates.

Thorough spray coverage of soybean plants is essential for optimum control. Utilize spray application techniques including sufficient water carrier per acre, pressure and proper nozzle

selection that ensure thorough coverage. **Caramba** should be applied in no less than 5 gallons of spray for aerial applications.

**Caramba** should be utilized as a part of a comprehensive Asian soybean rust control program that includes monitoring of the field and surrounding areas, and fungicide applications.

The key to adequate season long control of Asian soybean rust is careful monitoring and scouting of soybean fields all season, especially from initiation of flowering through pod fill. After the first application, maintain a thorough monitoring and scouting program.

**Caramba** provides both preventative and curative control of soybean rust. However, if symptoms or soybean rust lesions and/or pustules are present on soybean plants, some yield loss may have already occurred. If infection level of Asian soybean rust reaches or exceeds 3-5%, only limited control of Asian soybean rust may be achieved with any program. (Infection level = number of leaves with symptoms/signs of Asian soybean rust per 100 leaves).

### Conditions of Sale and Warranty

The Directions For Use of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Headline is a registered trademark of BASF.  
Caramba is a trademark of BASF.*

©2006 BASF Corporation  
All rights reserved.

007969-00000.20060324d.NVA 2006-04-267-0087

Supersedes NVA 2005-04-267-0161

BASF Corporation  
Agricultural Products  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company