



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Laredo[®] EC

Dow AgroSciences EPA Reg. No. 62719-412

Section 18 Quarantine Exemption for Control of Rust (*Phakopsora pachyrhizi*) in Soybean

(For Use and Distribution Only in the State of New York)

ATTENTION

- **Section 18 Quarantine Exemption:** This use is authorized under EPA quarantine exemption pursuant to Section 18 of the Federal Insecticide Fungicide and Rodenticide Act as amended.
- **This quarantine exemption is effective upon USDA confirmation that soybean rust is present in the continental U.S. and expires November 10, 2007.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Laredo[®] EC fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Laredo EC according to this Section 18 labeling is subject to all use precautions and limitations imposed by the label affixed to the product container for Laredo EC.

Directions for Use

Apply 4 to 8 fl oz of Laredo EC (0.0625 to 0.125 lb ai myclobutanil) per acre, using ground or aerial equipment, in an adequate spray volume to achieve good coverage and canopy penetration. For aerial application, apply Laredo EC in a minimum spray volume of 5 gallons per acre. For best results apply as early as possible, at first sign of disease, and make subsequent applications (up to 2) at 10- to 14-day intervals. For established disease, use the higher rate in the rate range.

Environmental Hazards

Groundwater: This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface water: This chemical can contaminate surface water through spray drift. Under some conditions, this chemical may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.


Specific Use Restrictions:

- **Do not** allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.
- **Do not** feed soybean forage and hay to livestock.
- **Preharvest Interval:** **Do not** make applications within 28 days of harvest.
- **Do not** make more than 2 applications of myclobutanil-containing products to soybeans per season.
- **Do not** apply more than 16 fl oz of Laredo EC (0.25 lb ai myclobutanil) to soybeans per acre per year.
- A maximum of 3 total applications using approved Section 18 products collectively are allowed under this soybean rust Section 18.
- **Rotation Crops** – Fields treated with myclobutanil (Laredo, Nova[®] and Rally[®]) can be rotated at any time to crops that are listed on a registered myclobutanil label immediately after the last treatment. **Do not** plant other crops within 30 days after the last application of a product containing myclobutanil.

ACCEPTED
MAY 19 2005

NEW YORK STATE DEPARTMENT
OF ENVIRONMENTAL CONSERVATION

Any adverse effects resulting from the use of Laredo EC under this quarantine exemption must be immediately reported to the New York Department of Agriculture.



®Trademark of Dow AgroSciences LLC

R153-048

File Code: 04-NY-15

Accepted: ___/___/___

Replaces R153-015.